


ENVIRONMENTAL AND OCCUPATIONAL EXPOSURE TO HEAVY METALS AND METALLOID AND DEVELOPMENT OF AMYOTROPHIC LATERAL SCLEROSIS: A POPULATION-BASED CASE CONTROL STUDY IN EMILIA-ROMAGNA AND SICILY

Federica Violi¹, Tommaso Filippini¹, Carlotta Malagoli^{1,2}, Maria Fiore³, Caterina Ledda³, Cristina Mauceri³, Angela Dimartino³, Jessica Mandrioli⁴, Nicola Fini⁴, Francesco Patti³, Margherita Ferrante³; Marco Vinceti^{1,2}



Affiliations:

- 1 CREAGEN - Environmental, Genetic and Nutritional Epidemiology Research Center, University of Modena and Reggio Emilia, Italy
- 2 Department of Neuroscience, University of Modena and Reggio Emilia, Modena, Italy
- 3 Department "G.F. Ingrassia", Section of Hygiene and Public Health and Laboratory of Environmental and Food Hygiene, University of Catania, Italy
- 4 Sant'Agostino Estense Hospital, Azienda Ospedaliero Universitaria Policlinico di Modena, Italy




Background

Amyotrophic lateral sclerosis (ALS) is a progressive, fatal neurodegenerative disease of the motor neuron. Its etiology is still largely unknown, except for some rare forms of genetic origin. Many environmental and occupational risk factors have been evaluated, including exposure to heavy metals, such as lead and mercury, and a metalloid, such as selenium. Clarifying the environmental component of ALS etiology could yield a better understanding of disease pathophysiology.

Methods

We performed a population based case-control study in three Italian provinces (Modena, Reggio Emilia and Catania) in order to assess the possible etiologic role of some environmental and occupational risk factors. We administered 877 questionnaires by mail or by person in a neurological office to newly-diagnosed ALS cases in the 2008-2011 period and age- and sex- matched population controls. The administered questionnaires collected information about personal, clinical and professional history, including job-related or hobby-related use of many trace elements and chemicals.



Results

Among the 162 returned questionnaires, 61 were cases and 101 controls, with an average response rate of 18.5%. The analyses, adjusted for age and sex, showed an increased risk when examining clinical information for reported occupational exposure to lead (odds ratio-OR 3.52, confidence interval-CI 95% 1.43 to 8.64, $p=0.006$), mercury (OR 6.16, 95% CI 0.66 to 57.72, $p=0.111$), while no increased risk was found to be associated with reported exposure to cadmium (OR 0.74, 95% CI 0.06 to 8.63, $p=0.814$). Finally, a slight although imprecise increase risk was found with occupational exposure to selenium (OR 1.67, 95% CI 0.22 to 12.39, $p=0.610$).

Conclusions

Our results seem to suggest a potential etiologic role of investigated trace elements and risk of ALS. However, the risk of possible selection bias due to the low response rate and information bias (i.e. recall bias) need to be carefully considered and worthy of further studies.

Bibliography

- Bozzoni V. et al, Amyotrophic lateral sclerosis and environmental factors. *Funct Neurol* 31, 7-19, 2016.
- Johnson FO et al., The role of environmental mercury, lead and pesticide exposure in development of amyotrophic lateral sclerosis, *Neurotoxicology*. 2009 Sep;30(5):761-5.
- Vinceti M et al, Lead, cadmium and mercury in cerebrospinal fluid and risk of amyotrophic lateral sclerosis: A case-control study, *J Trace Elem Med Biol*. 2017 Sep;43:121-125.

2017 SCIENTIFIC ANNUAL MEETING
ITALIAN ASSOCIATION FOR THE STUDY OF
TRACE ELEMENTS IN LIVING ORGANISMS – AISETOV

“Trace elements in human and animal health: focus on neurological disease”
Reggio Emilia - October 20, 2017

