

# The influence of outdoor air pollution on cadmium exposure assessment: a cross-sectional population-based Italian study

Filippini Tommaso<sup>1</sup>, Cherubini Andrea<sup>2</sup>, Maffei Giuseppe<sup>2</sup>, Greco Salvatore<sup>2</sup>, Malagoli Carlotta<sup>1</sup>, Malavolti Marcella<sup>1</sup>, Sieri Sabina<sup>3</sup>, Krogh Vittorio<sup>3</sup>, Vescovi Luciano<sup>4</sup>, Modenesi Marina<sup>4</sup>, Michalke Bernhard<sup>5</sup>, Vinceti Marco<sup>1</sup>

<sup>1</sup>Environmental, Genetic and Nutritional Epidemiology Research Center (CREAGEN), University of Modena and Reggio Emilia, Reggio Emilia, Italy;  
<sup>2</sup>Italy TerrAria srl, Milan, Italy;  
<sup>3</sup>Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei Tumori Milan, Italy;  
<sup>4</sup>IREN, Reggio Emilia and Piacenza, Italy;  
<sup>5</sup>Research Unit Analytical BioGeoChemistry, Helmholtz Zentrum München – German Research Center for Environmental Health GmbH, Munich, Germany.

## Background and Aims

Cadmium (Cd) poses serious environmental health hazards to humans. Cigarette smoking and diet are usually main sources of exposure in non-occupationally exposed subjects, while non-ferrous metal industrial production, fossil fuel combustion, cement


production and waste incineration are main anthropogenic sources of outdoor air Cd. The study aim was to assess the influence of outdoor air pollution on serum Cd levels in an Italian population.

## Methods

Outdoor exposure to particulate matter $\leq 10\mu\text{m}$  (PM<sub>10</sub>) from motorized traffic was assessed for fifty residents randomly-selected from Modena municipality. We geocoded the current residence of these subjects and modeled the corresponding ambient air concentration of particulate matter  $\leq 10\mu\text{m}$  (PM<sub>10</sub>) using the California LINE Source Dispersion Model version 4 (CALINE-4) as a proxy of environmental air Cd level. We compared these estimates with serum Cd, measured with inductively coupled plasma mass spectrometry. Information on smoking habits and Cd dietary intake were collected with a semi-quantitative food frequency questionnaire. We determined with both crude and multivariate linear regression models the influence of outdoor PM<sub>10</sub> levels, smoking and dietary Cd intake on serum Cd, computing  $\beta$ -coefficients and their 95% confidence interval (CI).




## Results

Median values (25th–75th) for serum and dietary Cd were 40.60 ng/l (30.05–53.50) and 13.36  $\mu\text{g}/\text{die}$  (10.45–16.77). Crude  $\beta$ -coefficients for PM<sub>10</sub>, dietary Cd and smoking on serum Cd levels were 0.617 (95% CI -0.194–1.428, P=0.133), 0.026 (-0.827–0.829, P=0.952) and 6.962 (-0.022–13.945, P=0.051), respectively. Corresponding adjusted values were 0.463 (-0.365–1.292, P=0.266), -0.036 (-0.866–0.793, P=0.930) and 6.057 (-1.175–13.289, P=0.099), respectively.



Modena Municipality Map with layers of interest for outdoor PM<sub>10</sub> exposure assessment

Sources of Cd	Crude			Adjusted		
	$\beta$	95% CI	P	$\beta$	95% CI	P
Outdoor PM <sub>10</sub> levels	0.632	(-0.275 – 1.539)	0.168	-0.473	(-0.451 – 1.399)	0.308
Dietary intake	0.026	(-0.827 – 0.879)	0.952	-0.050	(-0.884 – 0.785)	0.905
Smoking habits	6.962	(-0.022 – 13.945)	0.051	-6.199	(-1.020 – 13.419)	0.091



Figures and Table: linear regression analysis between serum cadmium (ng/L) and PM<sub>10</sub> ( $\mu\text{g}/\text{m}^3$ ), dietary intake ( $\mu\text{g}/\text{die}$ ) and smoking habits (smoking categorized as 0=never smokers, 1=former smokers, 2=current smokers). Multivariate model included as adjusting variable each factor alternatively.

## Conclusions

In our population, the most important factor influencing Cd serum content appears to be cigarette smoking, followed by outdoor air pollution (measured by PM<sub>10</sub> levels) and lastly diet, possibly for the limitations of dietary assessment methodology. In addition, other unmeasured factors could have influenced serum Cd content, such as a slow release from liver and kidney due to antecedent long-term exposure.

## Bibliography

- Vahter M, Berglund M, Nermell B, Akesson A. Bioavailability of cadmium from shellfish and mixed diet in women. *Toxicol Appl Pharmacol*. 1996;136:332-341.
- Nordberg M. Environmental exposure and preventive measures in Sweden and EU. *Biometals*. 2004;17:589-592.
- Druifuca A, Battaiotto S, Bengo I, Rossi D, Torriani L. Variante Generale al Piano Territoriale di Coordinamento Provinciale. Procedura di simulazione della mobilità delle persone. Milano: Provincia di Modena–Polinomia; 2007.
- Jarup L, Akesson A. Current status of cadmium as an environmental health problem. *Toxicol Appl Pharmacol*. 2009;238:201-208.



UNIMORE  
UNIVERSITÀ DEGLI STUDI DI  
MODENA E REGGIO EMILIA

Mail to Prof. Marco Vinceti  
at CREAGEN – Environmental, Genetic  
and Nutritional Epidemiology Research  
Center, University of Modena and Reggio  
Emilia, Via Campi 287 – 41125 Modena.  
marco.vinceti@unimore.it

Creagen  
Environmental, Genetic and Nutritional  
Epidemiology Research Center

fossil fuel heavy metal contaminants keywords  
main roads population PM<sub>10</sub> CALINE  
age spatial resolution smoking questionnaire assessment  
diet exposure methodology sample toxic  
limitations age resolution smoking cadmium exposure  
estimates regression environment air pollution outdoor indoor  
subjects waste incinerator highway foodstuff GIS epidemiology research correlation